



(11) **EP 1 459 763 A1**

(12) **EUROPEAN PATENT APPLICATION**  
published in accordance with Art. 158(3) EPC

(43) Date of publication:  
22.09.2004 Bulletin 2004/39

(21) Application number: 02781819.4

(22) Date of filing: 20.11.2002

(51) Int Cl.<sup>7</sup>: **A61K 45/00, A61K 31/131,  
A61K 31/4015, A61K 45/06,  
A61L 27/54, A61P 9/00,  
A61P 43/00**

(86) International application number:  
**PCT/JP2002/012091**

(87) International publication number:  
**WO 2003/043656 (30.05.2003 Gazette 2003/22)**

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR  
IE IT LI LU MC NL PT SE SK TR**  
Designated Extension States:  
**AL LT LV MK RO SI**

(30) Priority: 20.11.2001 JP 2001354931  
12.11.2002 JP 2002328138

(71) Applicants:  
• **Nipro Corporation**  
Kita-ku, Osaka 531-8510 (JP)

• **K.I.C. Co., Ltd.**  
Yokohama-shi, Kanagawa 236-0032 (JP)


(72) Inventor: **IKARI, Yuji**  
Suginami-ku, Tokyo 167-0041 (JP)

(74) Representative: **Merrifield, Sarah Elizabeth et al**  
**Boulton Wade Tennant**  
Verulam Gardens  
70 Gray's Inn Road  
London WC1X 8BT (GB)

(54) **DRUG FOR PREVENTING VASCULAR RESTENOSIS AND INSTRUMENT TO BE EMBEDDED  
IN VESSEL COATED WITH THE DRUG**

(57) A drug for preventing vascular restenosis comprises a compound that inactivates  $\alpha$ 1-protease inhibitor and  $\alpha$ 2-macroglobulin as active constituents, which are associated with occurrence of vascular restenosis after coronary artery intervention. Preferably, the drug is coated on a surface of an implantable intravascular device and locally administered to the affected area by application of the implantable intravascular device to a narrowed area.

Fig. 5



EP 1 459 763 A1

**Description****Technical Field**

5 [0001] The present invention relates to a vascular restenosis preventing agent and an implantable intravascular device coated therewith. More particularly, the invention relates to a drug for preventing vascular restenosis after coronary intervention, and an implantable intravascular device coated therewith.

**Background Art**

10 [0002] In late years, as a therapy for ischemic heart diseases resulting from coronary artery stenosis such as angina pectoris and myocardial infarction, a coronary intervention has been popularized rapidly in place of highly invasive coronary artery bypass graft surgery.

15 [0003] Typical coronary artery intervention is a percutaneous transluminal coronary angioplasty (PTCA) in which a lumen of a blood vessel is expanded by inserting a catheter with an inflatable balloon at a distal end thereof and then inflating the balloon at a stenosed area of a coronary artery. However, the therapy only by PTCA provides postoperative restenosis at a high frequency of about 40%, which has need of another therapy, causing the patient great mental and physical sufferings.

20 [0004] In order to solve such problems, a procedure, called stent placement, of placing a stent or a coiled or meshed cylindrical support of a metal or plastic at a widened stenosed area, has widely been brought into effect. This procedure has reduced the vascular restenosis rate to the halves.

[0005] The stent placement enables to prevent elastic recoil and vascular remodeling, but it is difficult to completely prevent the patient from vascular restenosis in the chronic stage. In fact, the stent placement provides about 20% probabilities of occurrence of an In-Stent Restenosis (ISR) after treatment. In particular, diabetic patients have higher frequencies of ISR.

25 [0006] It is conceived that the occurrence of ISR results from inflammation of the vessel walls caused by stent replacement, which causes intimal hyperplasia resulting from migration of smooth muscle cells from media to the intima.

[0007] For that reason, recent studies of the prevention of ISR have focused on inhibition of the neointimal hyperplasia, suggesting usefulness of drugs such as tranilast, paclitaxel, probucol, cilostazol and rapamycin. Some therapies employing stents coated with such a drug have been reported in Japanese unexamined patent publications such as, for example, JP2000-95706 (cf. page 3) and JP2000-249709 (cf. page 3). However, there is no generally accepted view that these drugs have remarkable effects on the prevention of ISR.

30 [0008] On the other hand, radiotherapies irradiating the coronary artery disease sites with  $\beta$ -rays or  $\gamma$ -rays to inhibit the neointimal hyperplasia have been put into practical use with good results. However, the radiotherapy has not yet been popularized since it causes some complications such as edge effects or late thrombosis and is difficult to control an exposed dose and radiation source.

35 [0009] In the current status, the prevention of vascular restenosis after coronary artery intervention is waiting for development of drugs or medical devices, which have fully satisfying effects on prevention of vascular restenosis after coronary artery intervention.

**Disclosure of Invention**


40 [0010] The inventors have studied solution of the above problems and found that the mechanism of ISR begins with migration of smooth muscle cells to fibrin adhered to the stent surfaces, and that  $\alpha_1$ -protease inhibitor,  $\alpha_1$ -antichymotrypsin and  $\alpha_2$ -macroglobulin in serum act as significant related factors in the initial stage where the fibrins adhere to the stent surface. These studies are reported in papers (Yuji Ikari et al, JOURNAL OF BIOLOGICAL CHEMISTRY, Vol.275, Issue 17, 12799-12805, April, 2000; Yuji Ikari et al, JOURNAL OF BIOLOGICAL CHEMISTRY Vol.276, Issue 15, 11798-11803, April, 2001). At the time of announcement of the papers, it was not clear what factor is an important factor related to ISR and thus there is no conclusion on the ISR-related factors.

50 [0011] Farther studies based on the above findings have been made on relationship between ISR and the level of the above factors in the blood for the patients treated by the stenting procedure. As a result, the inventors have concluded that the  $\alpha_1$ -protease inhibitor and  $\alpha_2$ -macroglobulin are related to narrowing of the blood vessel, but  $\alpha_1$ -antichymotrypsin has no relationship to the narrowing. The inventors have further studied on the above finding and achieved the present invention on the finding that ISR can be prevented effectively by deactivating  $\alpha_1$ -protease inhibitor and  $\alpha_2$ -macroglobulin with a certain substance.


55 [0012] According to the present invention, there is provided a drug for preventing vascular restenosis, which comprises, as an active constituent, a compound that deactivates  $\alpha_1$ -protease inhibitor.

[0013] The above compound that deactivates  $\alpha_1$ -protease inhibitor is a compound, or a salt thereof, represented by


the general formula:




wherein X is halogen, R<sub>1</sub> is hydrogen or an alkaline metal, R<sub>2</sub> is hydrogen or arylsulfonyl which may be being substituted, R<sub>1</sub> and R<sub>2</sub> taken together, can form a group represented by the general formulas:



(II-1)




(II-2)



(II-3)

or



(II-4)

wherein R<sub>3</sub> is a divalent group or bond of linear or branched C<sub>1</sub> - C<sub>5</sub> hydrocarbons which may have one or more unsaturated bonds which may have one or more unsaturated bonds.

[0014] In the above general formula, halogen atom expressed by X includes, for example, fluorine atom, chlorine atom, bromine atom and iodine atom, and a preferred halogen atom is chlorine atom. Examples of the alkaline metal represented by R<sub>1</sub> are lithium, sodium and potassium, but a preferred alkaline metal is sodium.

[0015] The arylsulfonyl represented by R<sub>2</sub> may have 1-5 substituents, preferably, 1-3 substituents, which may be the same or different with one another, at substitutable positions thereof. The arylsulfonyl includes phenylsulfonyl, naphthylsulfonyl and the like. As preferred examples of the above substituents, there may be used, for example, C<sub>3</sub>-C<sub>10</sub> cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; C<sub>2</sub>-C<sub>10</sub> alkenyl such as vinyl, allyl, 2-methylallyl, 2-butenyl, 3-butenyl, 3-octenyl; C<sub>2</sub>-C<sub>10</sub> alkynyl such as ethynyl, 2-propynyl, 3-hexynyl; C<sub>3</sub>-C<sub>10</sub> cycloalkenyl such as cyclopropenyl, cyclopentenyl and cyclohexenyl; C<sub>6</sub>-C<sub>10</sub> aryl such as phenyl and naphthyl; C<sub>7</sub>-C<sub>11</sub> aralkyl such as benzyl and phenylethyl; nitro; nitroso; hydroxyl; mercapto; cyano; oxo; thio; carbamoyl; mono or di- C<sub>1</sub>-C<sub>6</sub> alkylcarbamoyl such as methylcarbamoyl, dimethylcarbamoyl; C<sub>6</sub>-C<sub>14</sub> arylcarbamoyl such as phenylcarbamoyl; carboxyl; C<sub>1</sub>-C<sub>4</sub> alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl; C<sub>6</sub>-C<sub>14</sub> aryloxy-carbonyl such as phenoxycarbonyl; sulfo; halogen such as fluorine, chlorine, bromine and iodine; C<sub>1</sub>-C<sub>4</sub> alkoxy such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy; C<sub>6</sub>-C<sub>10</sub> aryloxy such as phenoxy; C<sub>1</sub>-C<sub>4</sub> alkylthio such as methylthio, ethylthio, propylthio, isopropylthio, butylthio, isobutylthio, sec-butylthio, tert-butylthio; C<sub>6</sub>-C<sub>10</sub> arylthio such as phenylthio; C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl such as methylsulfinyl, ethylsulfinyl, propylsulfinyl, isopropylsulfinyl, butylsulfinyl, isobutylsulfinyl, sec-butylsulfinyl, tert-butylsulfinyl; C<sub>6</sub>-C<sub>10</sub> arylsulfinyl such as phenylsulfinyl; C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl such as methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, butylsulfonyl, isobutylsulfonyl, sec-butylsulfonyl, tert-butylsulfonyl; C<sub>6</sub>-C<sub>10</sub> arylsulfonyl such as phenylsulfonyl; C<sub>1</sub>-C<sub>4</sub> alkoxy-sulfonyl such as methoxysulfonyl, ethoxysulfonyl, propoxysulfonyl, isopropoxysulfonyl, butoxysulfonyl, isobutoxysulfonyl, sec-butoxysulfonyl, tert-butoxysulfonyl; C<sub>6</sub>-C<sub>10</sub> aryloxy-sulfonyl such as phenoxysulfonyl; amino; C<sub>1</sub>-C<sub>11</sub> carboxyacylamino such as acetylamino, propionylamino, benzoylamino; mono- or di- C<sub>1</sub>-C<sub>4</sub> alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, dimethylamino, diethylamino; C<sub>3</sub>-C<sub>6</sub> cycloalkylamino such as cyclohexylamino; C<sub>6</sub>-C<sub>10</sub> arylamino such as anilino; trisubstituted silyl such as trimethylsilyl, tertbutyldimethylsilyl, triphenylsilyl, tert-butyl methoxyphenylsilyl; C<sub>1</sub>-C<sub>11</sub> carboxylic acyl such

as formyl, acetyl, benzoyl; a group selected from 3- to 6-membered heterocyclic groups containing 1-5 hetero atoms such as oxygen, sulfur or nitrogen, and condensed rings thereof, for example, such as 2- or 3-thienyl, 2- or 3-furyl, 1-, 2- or 3-pyrrolyl, 2-, 3- or 4-pyridyl, 2-, 4- or 5-oxazolyl, 2-, 4- or 5-thiazolyl, 1-, 3-, 4- or 5-pyrazolyl, 1-, 2-, 4- or 5-imidazolyl, 3-, 4- or 5-isoxazolyl, 3-, 4- or 5-isothiazolyl, 1,2,3- or 1,2,4-triazolyl, 2-, 4- or 5-pyrimidinyl, benzothiazolyl, benzoxazolyl, triazinyl, oxilanyl, aziridinyl, pyrrolidinyl, piperidinyl, morpholinyl, benzimidazolyl, quinolyl, isoquinolyl; an amino acid residue such as, for example, alanyl,  $\beta$ -alanyl, arginyl, cystathionyl, cystyl, glycyl, histidyl, homoseryl, isoleucyl, lanthionyl, leucyl, lysyl, methionyl, norleucyl, norvalyl, ornithyl, prolyl, sarcosyl, seryl, threonyl, thyronyl, tyrosyl, varyl, etc.

[0016] The above substituent may also have one or more substituents that may be the same or different from one another in the same or different substitutable positions. In that case, the further substituents are equivalent to the aforesaid substituents. If two or more substituents are used, their two substituents taken together, may form a divalent group such as, for example, C<sub>1</sub>-C<sub>6</sub> alkylene (e.g., methylene, ethylene, trimethylene, tetramethylene, propenylene), 3-oxapentamethylene, vinylene, benzylidene, methylenedioxy, 2-thiatrimethylene, oxalyl, malonyl, succinyl, maleoyl, phthaloyl, oxygen, sulfur, imino, azo, hydrazo and the like. If these substituents are divalent groups such as, for example, aryl, aralkyl, cycloalkyl, cycloalkenyl, aryloxy, arylthio, arylsulfinyl, arylsulfonyl, arylcarbamoyl, aryloxycarbonyl, aryloxysulfonyl, arylamino; cycloalkylamino, carboxylic acyl, carboxylic acylamino, tri-substituted silyl, and heterocyclic groups, 1 to 5 atoms of the substituents may further be substituted by one or more groups selected from the group consisting of halogen such as fluorine, chlorine, bromine and iodine; hydroxyl; nitro; cyano; C<sub>1</sub>-C<sub>4</sub> alkyl such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and tert-butyl; C<sub>2</sub>-C<sub>4</sub> alkenyl such as vinyl and allyl; C<sub>2</sub>-C<sub>4</sub> alkynyl such as, ethynyl, 2-propynyl and the like; phenyl; C<sub>1</sub>-C<sub>4</sub> alkoxy such as methoxy, ethoxy and the like; phenoxy; C<sub>1</sub>-C<sub>4</sub> alkylthio groups such as methylthio, ethylthio; and phenylthio group.

[0017] Further, if the substituents are groups selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylcarbamoyl, alkoxycarbonyl, alkoxysulfonyl, amino and alkylamino, 1 to 5 atoms of the substituent may further be substituted by one or more groups selected from the group consisting of halogen as mentioned above, hydroxyl, nitro; cyano; C<sub>1</sub>-C<sub>4</sub> alkoxy and C<sub>1</sub>-C<sub>4</sub> alkylthio groups.


[0018] As the divalent hydrocarbon group represented by R<sub>3</sub>, i.e., "the divalent group of linear or branched C<sub>1</sub>-C<sub>5</sub> hydrocarbons which may have one or more unsaturated bonds", there are C<sub>1</sub>-C<sub>5</sub> alkylene such as ethylene, trimethylene, tetramethylene, pentamethylene; C<sub>2</sub>-C<sub>5</sub> alkenylene such as vinylene, propenylene and the like.

[0019] In an embodiment of the present invention, the halogen atom in the above general formula (I) is chlorine atom.


[0020] In an embodiment of the present invention, the compound that deactivates  $\alpha_1$ -protease inhibitor is at least one of N-chlorosuccinimides.

[0021] According to the present invention, there is further provided a drug for preventing vascular restenosis, comprising as an active constituent, a compound that deactivates  $\alpha_2$ -macroglobulin.


[0022] As the above compound that deactivates  $\alpha_2$ -macroglobulin, it is preferred to use a compound or a salt thereof, represented by the general formula:



or



wherein R<sub>4</sub> is hydrogen or amino, R<sub>5</sub> is hydrogen, C<sub>1</sub>-C<sub>5</sub> alkyl that may be being substituted or carbamoyl that may be being substituted, p is an integer ranging from 1 to 3, Y is an acid radical, R<sub>4</sub> and R<sub>5</sub> taken together, may form a group represented by the general formula:



wherein m is an integer ranging from 1 to 6, in that case the  $\alpha_2$ -macroglobulin-deactivating compound may be being polymerized by ring-opening polymerization as represented by the general formula (V):



wherein n is an integer ranging from 10 to 2500.

[0023] The amino group represented by  $R_4$  in the above general formula includes, for example, a primary amino group, a secondary group or a tertiary amino group.

[0024] The alkyl in unsubstituted or substituted alkyl represented by  $R_5$  includes, for example,  $C_1$ - $C_5$  alkyl such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl and pentyl.

[0025] The alkyl or carbamoyl represented by  $R_5$  may have one or more substituents, which may be the same or different with one another, in substitutable positions thereof. As preferred examples of the above substituents, there may be used, for example,  $C_3$ - $C_{10}$  cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl;  $C_2$ - $C_{10}$  alkenyl such as vinyl, allyl, 2-methylallyl, 2-butenyl, 3-butenyl, 3-octenyl;  $C_2$ - $C_{10}$  alkynyl such as ethynyl, 2-propynyl, 3-hexynyl;  $C_3$ - $C_{10}$  cycloalkenyl such as cyclopropenyl, cyclobutenyl, cyclopentenyl and cyclohexenyl;  $C_6$ - $C_{10}$  aryl such as phenyl and naphthyl;  $C_7$ - $C_{11}$  aralkyl such as benzyl and phenylethyl; nitro; nitroso; hydroxyl; mercapto; cyano; oxo; thio; carbamoyl; mono or di-  $C_1$ - $C_6$  alkylcarbamoyl such as methylcarbamoyl, dimethylcarbamoyl;  $C_6$ - $C_{14}$  arylcarbamoyl such as phenylcarbamoyl; carboxyl;  $C_1$ - $C_4$  alkoxy-carbonyl such as methoxycarbonyl, ethoxycarbonyl;  $C_6$ - $C_{14}$  aryloxy-carbonyl such as phenoxycarbonyl; sulfo; halogen such as fluorine, chlorine, bromine and iodine;  $C_1$ - $C_4$  alkoxy such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy;  $C_6$ - $C_{10}$  aryloxy such as phenoxy;  $C_1$ - $C_4$  alkylthio such as methylthio, ethylthio, propylthio, isopropylthio, butylthio, isobutylthio, sec-butylthio, tert-butylthio;  $C_6$ - $C_{10}$  arylthio such as phenylthio;  $C_1$ - $C_4$  alkylsulfinyl such as methylsulfinyl, ethylsulfinyl, propylsulfinyl, isopropylsulfinyl, butylsulfinyl, isobutylsulfinyl, sec-butylsulfinyl, tert-butylsulfinyl;  $C_6$ - $C_{10}$  arylsulfinyl such as phenylsulfinyl;  $C_1$ - $C_4$  alkylsulfonyl such as methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, butylsulfonyl, isobutylsulfonyl, sec-butylsulfonyl, tert-butylsulfonyl;  $C_6$ - $C_{10}$  arylsulfonyl such as phenylsulfonyl;  $C_1$ - $C_4$  alkoxy-sulfonyl such as methoxysulfonyl, ethoxysulfonyl, propoxysulfonyl, isopropoxy-sulfonyl, butoxysulfonyl, isobutoxy-sulfonyl, sec-butoxy-sulfonyl, tert-butoxy-sulfonyl;  $C_6$ - $C_{10}$  aryloxy-sulfonyl such as phenoxysulfonyl; amino;  $C_1$ - $C_{11}$  carboxyacylamino such as acetylamino, propionylamino, benzoylamino; mono- or di-  $C_1$ - $C_4$  alkylamino such as methylamino, ethylamino, propylamino, isopropylamino, butylamino, dimethylamino, diethylamino;  $C_3$ - $C_6$  cycloalkylamino such as cyclohexylamino;  $C_6$ - $C_{10}$  arylamino such as anilino; trisubstituted silyl such as trimethylsilyl, tertbutyldimethylsilyl, triphenylsilyl, tert-butyl methoxyphenylsilyl;  $C_1$ - $C_{11}$  carboxylic acyl such as formyl, acetyl, benzoyl; a group selected from 3- to 6-membered heterocyclic groups containing 1-5 hetero atoms such as oxygen, sulfur or nitrogen, and condensed rings thereof, for example, such as 2- or 3-thienyl, 2- or 3-furyl, 1-, 2- or 3-pyrrolyl, 2-, 3- or 4-pyridyl, 2-, 4- or 5-oxazolyl, 2-, 4- or 5-thiazolyl, 1-, 3-, 4- or 5-pyrazolyl, 1-, 2-, 4- or 5-imidazolyl, 3-, 4- or 5-isoxazolyl, 3-, 4- or 5-isothiazolyl, 1,2,3- or 1,2,4-triazolyl, 2-, 4- or 5-pyrimidinyl, benzothiazolyl, benzoxazolyl, triazinyl, oxilanyl, aziridinyl, pyrrolidinyl, piperidinyl, morpholinyl, benzimidazolyl, quinolyl, isoquinolyl; an amino acid residue such as, for example, alanyl,  $\beta$ -alanyl, arginyl, cystathionyl, cystyl, glycyl, histidyl, homoseryl, isoleucyl, lanthionyl, leucyl, lysyl, methionyl, norleucyl, norvalyl, ornithyl, prolyl, sarcosyl, seryl, threonyl, thyronyl, tyrosyl, valyl, etc.

[0026] The above substituent may also have one or more substituents that may be the same with or different from the other, in the same or different substitutable positions. In that case, the further substituents are equivalent to the aforesaid substituents. If two or more substituents are used, their two substituents taken together, may form a divalent group such as, for example,  $C_1$ - $C_6$  alkylene (e.g., methylene, ethylene, trimethylene, tetramethylene, propenylene), 3-oxapentamethylene, vinylene, benzylidene, methylenedioxy, 2-thiatrimethylene, oxalyl, malonyl, succinyl, maleoyl, phthaloyl, oxygen, sulfur, imino, azo, hydrazo and the like. If these substituents are divalent groups such as, for example, aryl, aralkyl, cycloalkyl, cycloalkenyl, aryloxy, arylthio, arylsulfinyl, arylsulfonyl, arylcarbamoyl, aryloxy-carbonyl, aryloxy-sulfonyl, arylamino, cycloalkylamino, carboxylic acyl, carboxylic acylamino, tri-substituted silyl, and heterocyclic groups, 1 to 5 atoms of the substituents may further be substituted by one or more groups selected from the group consisting of halogen such as fluorine, chlorine, bromine and iodine; hydroxyl; nitro; cyano;  $C_1$ - $C_4$  alkyl such as methyl,

ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and tert-butyl; C<sub>2</sub>-C<sub>4</sub> alkenyl such as vinyl and allyl; C<sub>2</sub>-C<sub>4</sub> alkynyl such as, ethynyl, 2-propynyl and the like; phenyl; C<sub>1</sub>-C<sub>4</sub> alkoxy such as methoxy, ethoxy and the like; phenoxy; C<sub>1</sub>-C<sub>4</sub> alkylthio groups such as methylthio, ethylthio; and phenylthio group.

[0027] Further, if the substituents are groups selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylcarbamoyl, alkoxycarbonyl, alkoxysulfonyl, amino and alkylamino, 1 to 5 atoms of the substituent may further be substituted by one or more groups selected from the group consisting of halogen as mentioned above, hydroxyl, nitro; cyano; C<sub>1</sub>-C<sub>4</sub> alkoxy and C<sub>1</sub>-C<sub>4</sub> alkylthio groups.

[0028] The acid radical represented by Y includes chloro, carbonate, sulfate, nitrate and borate groups. Preferred acid radical is sulfate.

[0029] In a preferred embodiment of the present invention, the compound that deactivates  $\alpha_2$ -macroglobulin includes compounds containing an amino group. Such an amino-containing compound includes, for example, ammonia, amine compounds and salts thereof, urea, hydrazine and salts thereof, ammonium salts, imine and salts thereof, tobramycin, polylysines and lysozyme chloride, and the compound used is at least one compound selected from the group consisting of these compounds. When imine is used as the compound that deactivates  $\alpha_2$ -macroglobulin, it is preferred to use polyethylene imine having a molecular weight ranging from 500 to 100000, measured by gel filtration technique. In another embodiments, ammonium sulfate is employed as the compound that deactivates  $\alpha_2$ -macroglobulin.

[0030] The above compound that deactivates  $\alpha_2$ -macroglobulin may constitute a drug for prevention of vascular restenosis in combination with the above compound that deactivates  $\alpha_1$ -protease inhibitor.

[0031] In another aspect of the present invention, there is provided an implantable intravascular device coated with a drug for prevention of vascular restenosis that comprises, as an active constituent, a compound that deactivates  $\alpha_1$ -protease inhibitor and/or a compound that deactivates  $\alpha_2$ -macroglobulin. As an implantable intravascular device, it is preferred to use a stent.

#### Best Mode for Carrying Out the Invention

[0032] Firstly, the explanation will be made on a drug for prevention of vascular restenosis according to the present invention.


[0033] According to the present invention, there is no limit to the compound that deactivates  $\alpha_1$ -protease inhibitor, provided that the compound is able to deactivate the  $\alpha_1$ -protease inhibitor. Since an active center of the  $\alpha_1$ -protease inhibitor is considered to be blocked by an oxide, it is sufficient for the compound that deactivates  $\alpha_1$ -protease inhibitor to have a behavior to attack the oxide. Thus, it is preferred to use a compound having partial structure represented by formula:



Examples of the compounds having such a partial structure are cyclic imides or salts thereof derived from acyl of dibasic acids such as N-chloroxalimido, N-chloromalonimido, N-chlorosuccinimide, N-chlorophthalimide, N-chlormaleimide, N-chloroglutarimide; chloramine compounds obtained by replacing at least one hydrogen atom of ammonia with chlorine atom, or salts thereof. It is preferred to employ chloresuccinimide. These compounds may be used singly or in combination.

[0034] Further, there is no limit to the compound of the present invention that deactivates  $\alpha_2$ -macroglobulin, provided that the compound is able to deactivate  $\alpha_2$ -macroglobulin. Since it is considered that  $\alpha_2$ -macroglobulin is deactivated by the formation of a covalent bond between glutamic acid present in an active center of  $\alpha_2$ -macroglobulin and an amino group, it is preferred to use a compound having an amino group. As such a compound, there may be used ammonia; amine compounds or salts thereof having C<sub>1</sub>-C<sub>5</sub> alkyl replaced for one hydrogen atom of ammonia; urea having a carbamoyl group replaced for one hydrogen atom of ammonia; hydrazine or salts thereof having an amino group replaced for one hydrogen atom of ammonia; ammonium salts; and cyclic secondary amines having a divalent hydrocarbon replaced for two hydrogen atoms of ammonia, i.e., imines or salts thereof.

[0035] As the amine compound, it is preferred to use primary amine, in particular, methylamine or ethylamine, especially, methylamine. These amines may be used in the form of a salt thereof. As an ammonium salt, there may be used those such as, for example, ammonium hydrochloride, ammonium sulfate, ammonium carbonate, ammonium hydrogen carbonate, ammonium nitrate, ammonium borate. The most preferred ammonium salt is ammonium sulfate. The cyclic secondary amines having a divalent hydrocarbon replaced for two hydrogen atoms of ammonia includes imines represented by the general formula:



wherein k is an integer ranging from 1 to 6, such as, for example, ethylene imine, pyrrolidine and piperidine. A preferred imine is ethylene imine. The ethylene imine may be used in the form of polyethylene imine obtained by ring opening polymerization and expressed by the general formula:



wherein n is an integer ranging from 10 to 2500. In that case, preferred ethylene imine is polyethylene imine having a molecular weight ranging from 500 to 100000, measured by gel filtration technique. These drugs may be used singly or in combination with two or more drugs.

[0036] Further, if the C1-C5 alkyl with a substituent has any further substituent, examples of such a compounds include tobramycin, polylysine, elcatonin and lysozyme chloride.

[0037] The compound that deactivates  $\alpha_1$ -protease inhibitor and the compound that deactivates  $\alpha_2$ -macroglobulin may be administered in the form of bulk powder, but they are generally administered in the form of preparations prepared by mixing the component with any component selected from the group consisting of additives, fillers (e.g., calcium carbonate, lactose, crystallized cellulose, starches, porous materials, etc.), disintegrants (e.g., low-substituted hydroxypropylcellulose, carboxymethylcellulose calcium, etc.), binders (e.g., dextrin, gelatine, pullulans, etc.), lubricants (e.g., magnesium stearate, calcium stearate, etc.), coating agents (e.g., methyl cellulose, ethyl cellulose and the like), surfactants (e.g., polyoxyethylene hydrogenated castor oils, polysorbate, etc.), elasticizer (e.g., triethyl citrate, triacetin, etc.), antioxidant agents (e.g., citric acid,  $\alpha$ -tocopherol, etc.) and preservative (e.g., parabens and the like), and then making a suitable amount of the resultant mixture to a desired dosage form.

[0038] The dosage form includes, for example, tablets, balls, capsules, granules, subtle granules, powder, syrup, injectable solutions, suspensions, fat emulsion preparations, inhalants, ointment, etc., any of which may be prepared in the ordinary manner, for example, as described in Japanese Pharmacopoeia. Among them, it is preferred to use a dosage form suitable for parenteral and topical administration such as injectable solutions, suspensions and fat emulsion preparations.

[0039] A more preferred process is administration of a drug to an affected area directly with a drug delivery catheter or an implantable intravascular device coated with the drug. The most preferred process is placement of a stent coated with an drug for prevention of restenosis in the affected area. A process employing a stent is simple and easy in insertion, so that it requires no specialized technique and is short in time required for implantation of the stent in the affected area. Further, the stent coated with an drug for prevention of restenosis is locally implanted in the affected area, a dosage of the drug for regular adult is small and causes no side issues such as side effects.

[0040] Further, the dosage of drug for prevention of restenosis according to the present invention varies with administration routes, a patient's symptoms, age or weight. It is preferred to administer the drug such that a local concentration of the compound at the narrowed area of the vessel ranges from 1 to 100 mM. The concentration less than 1 mM does not have much effect on prevention of vascular restenosis, while the concentration exceeding 100 mM may cause side effects. The local concentration of the active component of the drug for prevention of vascular restenosis ranges from, preferably, 70 to 90 mM and, most preferably, is about 80 mM. To keep the local concentration, it is preferred to administer the drug for prevention of restenosis to the area parenterally and topically. This may be done, for example, by placement of an implantable intravascular device, especially a stent, coated with the drug for prevention of vascular restenosis of the present invention in the affected area, to achieve sustained-release of the drug.

[0041] Respective constituents and preparation process of the drug for prevention of vascular restenosis according to the present invention are described in detail in Japanese patent publications JP2000-249709(A), JP2000-95706(A) and Japanese Patent No. 3146009, which are hereby incorporated by reference.

[0042] Though the drug for prevention of vascular restenosis of the present invention has been explained as above, the vascular restenosis can be effectively prevented by use of a drug delivery balloon or a stent coated with the drug for prevention of vascular restenosis of the present invention to perform local administration of the drug.

[0043] Next, the explanation will be made on the implantable intravascular device of the present invention, making reference to a stent as an example.

[0044] A stent which is a preferred embodiment of implantable intravascular device of the present invention is characterized by the fact that a surface of the stent is coated with a drug for prevention of vascular restenosis. In general,

the stent has a cylindrical form, has a coiled body portion or a meshed body portion having plural slots. But the stent is not limited to these configurations and may take any configuration, provided that it can maintain a size of the lumen after widening the narrowed area of the vessel. Also, the stent may have an expansion mechanism of a self-expandable type or a balloon expandable type.

**[0045]** Suitable materials for the stent includes metals such as stainless steel, titanium, tantalum, aluminum, tungsten and nitinol; plastics such as polyamides, polyethylene, polypropylene and hardly-hydrolyzable polyether polyurethane. However, there is no limit to the material for the stent, provided that the material is biocompatible and has been proved as a material for medical use.

**[0046]** Examples of the stents includes Palmaz-Schatz (trademark) stent of Johnson & Johnson, MULTI-LINK (trademark) stent of Advanced cardiovascular systems, Cordis (trademark) stent of Cordis Corporation and Microstent-GFX (trademark) of Medotronic Arterial Vascular Engineering, Inc., which can be used with good results.

**[0047]** The stent of the present invention may be obtained by applying a coating mixture containing the drug for prevention of vascular restenosis of the present invention to the above stent in the conventionally known manner. Methods for coating to stents are described in Japanese patent unexamined publications JP2000-350786 A and JP2000-51367A, which are hereby incorporated by reference. However, these coating methods can be applied to implantable intravascular devices other than stent. The following is explanation of a method for coating a drug for prevention of vascular restenosis to the stent.

**[0048]** The drug-coated stent may be obtained by (1) a method comprising steps of mixing a drug for prevention of vascular restenosis with any one of various polymers to prepare a coating mixture, and then spraying the resultant mixture on a stent or dipping the stent into the resultant mixture, or (2) a method comprising steps of providing a hydrophilic coating of a polymer on surface of a stent and then spraying the drug for prevention of vascular restenosis on a stent or dipping the stent into the drug for prevention of vascular restenosis. The drug for prevention of vascular restenosis is used after formulation to a suitable dosage form such as a solution, powder or the like, to make it easy to mix the drug with any one of the various polymers. If a powder type drug for prevention of vascular restenosis is mixed with polymer, it is necessary to sufficiently reduce granularity (primary particles and secondary particles such as aggregates or lumps) to provide the stent with smooth surfaces after coating as well as to prevent slots of the stent from being clogged with the drug.

**[0049]** The above coating mixture may includes one or more additives such as, for example, diluents, carriers, fillers, stabilizing agents and the like. For example, one or more hydrophilic polymers selected from biocompatible hydrophilic polymers may be added to a biocompatible hydrophobic coating agent to modify release profile of the drug. Also, one or more hydrophobic polymers may be added to a biocompatible hydrophilic coating agent to modify release profile of the drug. For example, the release profile may be modified by incorporating a hydrophilic polymer selected from the group consisting of polyethylene oxide, polyvinyl pyrrolidone, polyethylene glycols, carboxymethylcellulose, hydroxymethylcellulose and combination thereof into an aliphatic polyester coating agent. An appropriate amount of the hydrophilic polymer added to the hydrophobic coating agent or that of hydrophobic polymer added to the hydrophilic coating agent is determined according to the release profile of the drug for prevention of vascular restenosis in vitro or in vivo.

**[0050]** There is no limit to the polymers used in the present invention if they are biocompatible. Examples of the polymers are polyamides, polyolefines (including polypropylene, polyethylene), nonabsorbable polyesters (including polyethylene terephthalate and the like) and biologically absorbable aliphatic polyesters (lactic acid, glycolic acid, lactide, glycolide, *p*-dioxane, trimethylenecarbonates,  $\epsilon$ -caprolactam, homopolymers thereof and copolymers derived from a mixture of these materials).

**[0051]** Further, the polymer may be mixed with a highly flexible, high molecular elastomer to provide coating layers with flexibility. The elastomers that can be used include, for example, various elastomers of polystyrene, polyolefine, polyesters, polyamides, silicone, polyurethane, fluorinated elastomer, natural rubber, and copolymers or polymer alloys of these materials. Among them, the most preferred material is a segmented polyurethane polymer film with a high flexibility and a high strength. It is preferred to use segmented polyurethane polymer films including flexible polyether soft-segments and hard segments of aromatic and urea bond-rich portions, the soft segments and hard segments forming a microstructure based on phase separation. Such segmented polyurethane polymer films have sufficient strength and are excellent in antithrombogenicity. In other words, the segmented polyurethane polymer films are biocompatible and can be expanded sufficiently without causing rupture.

**[0052]** An amount of the drug for prevention of vascular restenosis to be coated on the stent varies with the administration route, a patient's symptoms, age or weight. The quantity consumed of the drug for prevention of vascular restenosis ranges from about 0.001 w/v% to about 70 w/v%, more generally, from about 0.001 w/v% to about 60 w/v% and, most generally, from about 0.001 w/v% to about 45 w/v% with respect to the total amount of the coating mixture.

**[0053]** According to the present invention, the stent explained above makes it possible to provide low invasive means for prevention of vascular restenosis in a short time after coronary interventions as well as to effectively prevent the vascular restenosis, thus making it possible to reduce mental and physical pains of the patient.



[0054] It is to be noted that the coronary interventions with possibility of postoperative vascular restenosis, to which the drug for prevention of vascular restenosis of the present invention and implantable intravascular device coated therewith are applied, include, without being limited thereto, Directional Coronary Atherectomy (DCA), Transluminal Extraction Catheter (TEC), rotating atherectomy coronary angioplasty (Rotablator) and Laser Angioplasty, in addition to the aforesaid PTCA and stent placement.

#### BRIEF DESCRIPTION OF DRAWINGS

[0055]

Fig. 1 is a graph illustrating relationship between decreasing index for vascular restenosis rate and concentrations of  $\alpha_1$ -protease inhibitor,  $\alpha_2$ -macroglobulin and  $\alpha_1$ -antichymotrypsin in serum;

Fig. 2 is a graph illustrating relationship between presence or non-presence of vascular restenosis and concentration of  $\alpha_2$ -macroglobulin in serum for both diabetic patient group and nondiabetic patient group;

Fig. 3 is a graph illustrating relationship between a concentration of N-chlorosuccinimide in fibrin and migration incidence of vascular plain muscles;

Fig. 4 is a graph illustrating Intima-Media Thickness (I/M ratio) for an N-chlorosuccinimide-treated group and a control group (right common iliac artery 3-times injury model);

Fig. 5 is a graph illustrating effects of methylamine, ethylamine, polyethylene imine with molecular weight of 600 to 10000, tobramycin, ammonia, lysozyme chloride and polylysine on deactivation of  $\alpha_2$ -macroglobulin.

#### EXAMPLES

[0056] The invention will be specifically explained below with reference to experimental tests.

[0057] In the examples, The test of significant difference on results for respective experimental tests was carried out by Pearson text with a SAS (trademark) system Ver. 6.12, made by SAS Institute Japan, unless otherwise specified. Symbols in experimental tests respectively denote the followings, unless otherwise stated. a1P1:  $\alpha_1$ -protease inhibitor, a2M:  $\alpha_2$ -macroglobulin, a1ACT:  $\alpha_1$ -antichymotrypsin, p: rate of rejection, R: correlation coefficient, M: mol concentration (mol/L), mM: millimol concentration (mmol/L), mg: milligram, w/v%: weight/volume percentage, ISR: In-stent restenosis.

[0058] The followings are explicative of experimental results on case patients received stenting, which reveals that  $\alpha_1$ -protease inhibitor and  $\alpha_2$ -macroglobulin are ISR related factors as mentioned below.

[EXPERIMENT 1] Influences of concentrations of  $\alpha_1$ -protease inhibitor and  $\alpha_2$ -macroglobulin in serum on reduction of vascular lumen diameter

[0059] The objects of experiments were case patients with coronary artery diseases requiring coronary-artery intervention, who are de novo lesion and have received stent placements.

[0060] In order to prevent the patients from acute and subacute thrombotic occlusion, pretreatments were performed by administering aspirin (81 mg or 162 mg) and ticlopidine (200 mg) or cilostazole (200 mg) for 7 days prior to the therapy. There is no limit to the length of lesion and the selection of a stent was left to the discretion of the operator. Stenting was succeeded in 91 cases and 110 lesions. Among them, the 107 lesions were subjected to follow-up coronary angiography after 6 months. These 107 lesions were tested for verifications of relationship between concentrations of  $\alpha_1$ -protease inhibitor,  $\alpha_2$ -macroglobulin in blood and inner diameter of the vessel. The inner diameter of the blood vessel was measured before and after stent placement and after 6 months with a cardiovascular analysis software, QCA-CMS of CMS, and loss index which represents the level of vascular restenosis was determined from acute gain and late loss.

[0061] Blood serums were collected from the respective case patients after 24 hours of the therapy and subjected to the nephelometric analysis to determine concentrations of  $\alpha_1$ -protease inhibitor and  $\alpha_2$ -macroglobulin along with that of  $\alpha_1$ -antichymotrypsin as the control material. The measurements were carried out in an orderly manner by using measurement kits made by Dade Behring Marburg for  $\alpha_1$ -protease inhibitor and  $\alpha_2$ -macroglobulin and measurement kits made by Hoechst Japan for  $\alpha_1$ -antichymotrypsin.

[0062] Results are shown in Fig. 1. In Fig. 1, (A) denotes the relationship between the concentrations of  $\alpha_1$ -protease inhibitor in serum and loss index, (B) denotes the relationship between the concentrations of  $\alpha_2$ -macroglobulin in serum and loss index, and (C) denotes the relationship between the concentrations of  $\alpha_1$ -antichymotrypsin in serum and loss index. In each diagram, a vertical axis indicates the loss index, while a horizontal axis indicates the concentration of respective factors in serum.

[0063] As shown in Fig. 1, the mean values of concentrations of  $\alpha_1$ -protease inhibitor,  $\alpha_2$ -macroglobulin and  $\alpha_1$ -antichymotrypsin in serum were fallen in the normal range. As can be seen from the results shown in Fig. 1, there was

positive correlation between the loss index and  $\alpha_1$ -protease inhibitor or  $\alpha_2$ -macroglobulin, but there was no correlation between  $\alpha_1$ -antichymotrypsin and the loss index. In other words, it was confirmed that there is statistical correlation between  $\alpha_1$ -protease inhibitor or  $\alpha_2$ -macroglobulin and ISR.

[0064] Further, Fig. 2 shows the relationship between the concentration of  $\alpha_2$ -macroglobulin in serum and ISR, obtained by separating the case patients into a diabetic group and a nondiabetic group. In that figure, a black circle denotes the presence of ISR, while a white circle denotes no presence of ISR. The presence or absence of ISR was determined by measurement of the inner diameter of the blood vessel with a quantitative coronary angiography, and more than 50 % diameter stenosis was judged to be ISR.

[0065] As shown in Fig. 2, the concentrations of  $\alpha_2$ -macroglobulin in serum for the diabetic group are higher than that of the nondiabetic group. The data for the nondiabetic group have no correlation between concentration of  $\alpha_2$ -macroglobulin in serum and ISR. In contrast therewith, the data for the diabetic group have a clear correlation between concentration of  $\alpha_2$ -macroglobulin in serum and ISR since the increase in concentration of  $\alpha_2$ -macroglobulin in serum have a high tendency to increase the occurrence of ISR.

[0066] For example, in the case patients of the diabetic group, the concentration of  $\alpha_2$ -macroglobulin higher than the normal upper limit of  $0.35 \times 10^{-5}$  M causes a 75 % occurrence of ISR, while the concentration of  $\alpha_2$ -macroglobulin lower than  $0.35 \times 10^{-5}$  M provides a 0% occurrence of ISR. Thus, in the case patients of the diabetic group, it was revealed that the concentration of  $\alpha_2$ -macroglobulin in serum has a considerably strong correlation with the occurrence of ISR.

[0067] In other words, it can be said that the vascular restenosis preventing agent according to the present invention is a highly effective drug for diabetic patients.

[0068] Accordingly, it will be seen that the prevention of vascular restenosis can be expected by inhibiting  $\alpha_1$ -protease inhibitor and  $\alpha_2$ -macroglobulin.

[0069] Although serum factors associated with the vascular restenosis have been studied by investigators in the world, at the present moment there is no report on the serum factor that can be revealed to have such a correlation therewith.

[0070] The next is an explanation on test results showing pharmacologic effects of the compound (I).

#### [EXPERIMENT 2] Influences of N-chlorsuccinimide in fibrin on the migration frequencies of vascular smooth muscles

[0071] The vascular smooth muscles were collected from human vital arteries and established HNB18E6E7 was used as cell lines. The vascular smooth muscles were incubated in 6-well plates to produce confluent cultures. The resultant supernatant fluids were washed three time with warmed phosphate buffer solution (PBS) and softly added with 3 mg/mL of fibrinogen and 0.8 mL of Waymouth culture solution including 10 w/v% human serum. Then, the supernatant fluids were respectively added with N-chlorsuccinimide to adjust the final concentration of N-chlorsuccinimide to 0.08 mM, 0.8 mM, 8 mM and 80mM, and gelled by addition of thrombin. The resultant gelled supernatants were incubated at 37 ° C for 24 hours, and the numbers of the smooth muscle cells migrated into the gel was counted with a phase contrast microscope. For each concentration, the experiments were carried out to make n = 3. Results are shown in Fig. 3. The ordinate denotes the frequencies of migration of vascular smooth muscles into the gel, while the abscissa denotes the concentration of N-chlorsuccinimide.

[0072] As illustrated in Fig. 3, the frequency of migration of the vascular smooth muscles decreases apparently with increasing concentration of N-chlorsuccinimide, and the number of migrating vascular smooth muscles vanished completely at 80 mM. The results proved that N-chlorsuccinimide that is an inhibitor of  $\alpha_1$ -protease inhibitor has a beneficial effect on the prevention of vascular restenosis.

[0073] Although N-chlorsuccinimide was used in the above experiment, since the active centers of  $\alpha_1$ -protease inhibitor are protected by oxides, it is possible to use any other compound having chlorine atom and an oxidizing ability and including a partial structure represented by the formula:



[0074] Explanation will be made on experimental results proving medicinal values of the compound(I).

#### [EXPERIMENT 3] Influences of N-chlorsuccinimide on neointimal formation in vascular injury model

[0075] There were used a group of animals consisted of 5 (five) N2W male rabbits aged 2 months. Vascular injury models were prepared by introducing a Fogarty catheter, Model e-060-2F, made by Baxter corporation (that is an arterial embolectomy balloon catheter) into the right common iliac artery through the right femoral artery under pentobarbital anesthesia, and then abrading the right common iliac artery 3 times under conditions of the balloon being

inflated (balloon pressure: 0.5 - 1.0 atm). After injuring the vascular endothelium, hydrophilic balloons made by Boston Scientific Corporation, of an N-chlorosuccinimide-treated group prepared by immersing the balloons in 80 mM N-chlorosuccinimide, and of a control group prepared by immersing the balloons in PBS, were respectively inflated in the right common iliac arteries to locally administer the drug to each injured area.

[0076] After 4 weeks from the treatment, the arteries were surgically removed, fixed with 10w/v% neutral buffered formalin fixative and then sectioned to prepare arterial ring tissue sections. After preparing elastic fiber stained tissue samples by EVG (Elastic Van Gieson) staining, the samples were subjected to measurements of intimal area (I) and medial area (M) to determine their ratio (I/M) as the degree of intimal thickening. Results are shown in Fig. 4, in which the ordinate denotes the degree of intimal thickening (I/M ratio), and the abscissa denotes the treated group names.

[0077] As illustrated in Fig.4, In the group treated with N-chlorosuccinimide, the neointimal hyperplasia is considerably inhibited. It is proved that the compound is effective against the vascular restenosis.

[0078] Explanation will be made on experimental results proving medicinal values of the compound (III-1).

[EXPERIMENT 4] Effects of methylamine, ethylamine, polyethyleneimines with molecular weights of 600 and 10000, tobramycin, ammonia, lysozyme chloride and polylysine on deactivation of  $\alpha_2$ -macroglobulin

[0079] Utilizing the facts that trypsin is inhibited by  $\alpha_2$ -macroglobulin and that trypsin causes yellow coloring of a coloring matrix S2222 resulting from decomposition thereof, verification was carried out for effects of methylamine, ethylamine, polyethyleneimines with molecular weights of 600 and 10000, tobramycin, ammonia, lysozyme chloride and polylysine on deactivation of  $\alpha_2$ -macroglobulin. A concrete method will be given below.


[0080] Firstly, each of methylamine, ethylamine, polyethyleneimines with molecular weights of 600 and 10000, tobramycin, ammonia, lysozyme chloride and polylysine was diluted to 0.0001 w/v%, 0.001w/v%, 0.01 w/v%, 0.1 w/v% and 1 w/v% to prepare a series of diluted solutions as sample solutions. 50  $\mu$ L of each sample solution was added with 50  $\mu$ L of  $2.5 \times 10^{-4}$  w/v%  $\alpha_2$ -macroglobulin and left for 5 minutes at room temperature. Then, the sample solution was added with 50  $\mu$ L of a 5 w/v% trypsin EDTA solution, left for 5 minutes at room temperature, and then added with 10  $\mu$ L of the coloring matrix S2222. After leaving the resultant sample solution for 20 minutes at room temperature, the sample solution was added with 50  $\mu$ L of 10 w/v% acetic acid solution to stop reactions, and then subjected to measurement of absorption of light with a wave length of 405 nm, using 50 mM tris(hydroxymethyl)aminomethane buffer solution of pH 8.0 as the blank. The dilution of all the samples and agents was carried out by a 50 mM tris (hydroxymethyl)aminomethane buffer solution of pH 8.0. Results are shown in Fig. 5, in which the ordinate denotes the deactivation rate of  $\alpha_2$ -macroglobulin, and the abscissa denotes the concentrations of respective compounds.

[0081] As shown in Fig. 5, in each compound, increasing concentration thereof deactivates  $\alpha_2$ -macroglobulin. Polylysine and lysozyme chloride provide 100% deactivation rate of  $\alpha_2$ -macroglobulin at the concentration of 0.1 w/v%, while methylamine provides about 100% deactivation rate of  $\alpha_2$ -macroglobulin at the concentration of 1 w/v%. Further, polymethylene imine with molecular weights of 10000 provides about 60% deactivation rate of  $\alpha_2$ -macroglobulin at the concentration of 0.1 w/v%. Since each compound has the amino group, it is considered that  $\alpha_2$ -macroglobulin is deactivated by the formation of covalent bonds between the amino group and active centers of  $\alpha_2$ -macroglobulin.


[0082] The vascular restenosis preventing agent according to the present invention and the implantable intravascular device coated therewith deactivate  $\alpha_1$ -protease inhibitor and  $\alpha_2$ -macroglobulin, which are factors being closely related to generation of vascular restenosis, thus making it possible to effectively prevent and treat the restenosis of the blood vessel expanded by coronary artery intervention. The above restenosis preventing agents provide sufficient effects even at low concentrations, have very little side effects, and in particular work on diabetic patients effectively. In addition, in case that the above restenosis preventing agents is locally applied to the affected area with an implantable intravascular device, in particular, a stents coated with the agent, the effects of the agent increase greatly and a risk of side effects is considerably lowered.

#### Claims


1. A drug for preventing vascular restenosis, containing as an active constituent, a compound that inactivates  $\alpha_1$ -protease inhibitor.
2. The drug for preventing vascular restenosis according to claim 1, wherein the compound that inactivates  $\alpha_1$ -protease inhibitor is a compound, or a salt thereof, represented by the general formula:




wherein X is halogen, R<sub>1</sub> is hydrogen or an alkaline metal, R<sub>2</sub> is hydrogen or arylsulfonyl which may be being substituted, R<sub>1</sub> and R<sub>2</sub> taken together, can form a group represented by the general formula:




(II-1)



(II-2)




(II-3)




(II-4)

wherein R<sub>3</sub> is a divalent group of a linear or branched C<sub>1</sub>-C<sub>5</sub> hydrocarbon which may have one or more unsaturated bonds, or a divalent bonding group.

3. The drug for preventing vascular restenosis according to claim 2, wherein the halogen atom is a chlorine atom.
4. The drug for preventing vascular restenosis according to claim 1, wherein the compound that inactivates α<sub>1</sub>-protease inhibitor is N-chlorosuccinimide
5. An implantable intravascular device coated with a drug for preventing vascular restenosis defined in claim 1.
6. The implantable intravascular device according to claim 5, wherein said device is a stent.
7. A drug for preventing vascular restenosis, containing as an active constituent, a compound that inactivates α<sub>2</sub>-macroglobulin.
8. The drug for preventing vascular restenosis according to claim 7, wherein the compound that inactivates α<sub>2</sub>-macroglobulin is a compound or a salt thereof, said compound being represented by the general formula:




or




(III-2)

wherein  $\text{R}_4$  is hydrogen or amino,  $\text{R}_5$  is hydrogen,  $\text{C}_1$ - $\text{C}_5$  alkyl that may be being substituted or carbamoyl that may be being substituted,  $p$  is an integer ranging from 1 to 3,  $\text{Y}$  is an acid radical,  $\text{R}_4$  and  $\text{R}_5$  taken together, may form a group represented by the general formula:



wherein  $m$  is an integer ranging from 1 to 6, in that case the  $\alpha_2$ -macroglobulin-inactivating compound may be being polymerized by ring-opening polymerization as represented by the general formula (V):



(V)

wherein  $n$  is an integer ranging from 10 to 2500.

9. The drug for preventing vascular restenosis according to claim 8, wherein the compound that inactivates  $\alpha_2$ -macroglobulin is at least one compound selected from the group consisting of ammonia, methylamine, ethylamine, salts thereof, urea, tobramycin, polylysines and lysozyme chloride.

10. The drug for preventing vascular restenosis according to claim 8, wherein the compound that inactivates  $\alpha_2$ -macroglobulin is polyethylene imine having a molecular weight ranging from 500 to 100000, measured by gel filtration technique.

11. The drug for preventing vascular restenosis according to claim 8, wherein the compound that inactivates  $\alpha_2$ -macroglobulin is ammonium sulfate.

12. The drug for preventing vascular restenosis according to claim 7, further comprising a compound that inactivates  $\alpha_1$ -protease inhibitor in combination with said compound.

13. An implantable intravascular device coated with a drug for preventing vascular restenosis defined in claim 7 or 12.

14. The implantable intravascular device according to claim 13, wherein said device is a stent.

Fig. 1




Fig. 2




Fig. 3






Fig. 4





Fig. 5



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/12091

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>		
Int.Cl. <sup>7</sup> A61K45/00, 31/131, 31/4015, 45/06, A61L27/54, A61P9/00, 43/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols)		
Int.Cl. <sup>7</sup> A61K45/00, 31/131, 31/4015, 45/06, A61L27/54, A61P9/00, 43/00		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
MEDLINE (STN), CAPLUS (STN), BIOSIS (STN), EMBASE (STN), WPIDS (STN), REGISTRY (STN)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96/40098 A2 (NEORX CORP.), 19 December, 1996 (19.12.96), Abstract & US 5595722 A & AU 9662773 A & EP 833624 A2 & US 5770609 A & JP 11-510479 A & US 2002/0068731 A1 & US 2002/0182659 A1	7, 13, 14
X	WO 94/26303 A1 (NEORX CORP.), 24 November, 1994 (24.11.94), Abstract & US 5472985 A & EP 710116 A1 & US 5599844 A & JP 08-510451 A & US 5773479 A & US 5945456 A	7, 13, 14
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 19 February, 2003 (19.02.03)		Date of mailing of the international search report 11 March, 2003 (11.03.03)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1998)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/12091

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94/26291 A1 (NEORX CORP.), 24 November, 1994 (24.11.94), Abstract & US 5733925 A & EP 702557 A1 & US 5811447 A & JP 08-510209 A & US 6074659 A & US 2002/0013275 A1	7,13,14
Y	JP 2000-095706 A (Ajinomoto Co., Inc.), 04 April, 2000 (04.04.00), Par. No. [0005] (Family: none)	1-9,12,14
Y	IKARI Yuji, $\alpha$ 1-Proteinase Inhibitor, $\alpha$ 1-Antichymotrypsin, and $\alpha$ 2-Macroglobulin Are the Antiapoptotic Factors of Vascular Smooth Muscle Cells. The Journal of Biological Chemistry, 13 April, 2001 (13.04.01), Vol.276, No.15, pages 11798 to 11803, abstract	1,5-7,12-14
Y A	US 5057414 A (BEHRINGWERKE AG.), 15 October, 1991 (15.10.91), Whole article & EP 297597 A2 & DE 3722082 A1 & CA 1338681 A1 & AU 8818615 A1 & JP 64-030599 A	1-9,12,14 10
Y	DE 19712565 A1 (STIEF, Thomas W.), 01 October, 1998 (01.10.98), Claim 9 (Family: none)	1,2,5,6
Y	STIEF T. W. et al., Singlet oxygen inhibits agonist-induced P-selectin expression and formation of platelet aggregates. Chemical abstracts. Vol.136, 2001, No.241367, & Clinical and Applied Thrombosis/Hemostasis. 2001, Vol.7, No.3, pages 219 to 224	1,2,5,6
X	WO 01/45862 A1 (INNERDYNE, INC.), 28 June, 2001 (28.06.01), Abstract & US 2002/0009604 A1 & EP 1251973 A1	7-9,13,14
Y	JP 2001-213983 A (Kabushiki Kaisha Ion Kogaku Kenkyusho), 07 August, 2001 (07.08.01), Claim 4 (Family: none)	7-9,11,13, 14
A	WO 98/47948 A1 (CALIFORNIA INSTITUTE OF TECHNOLOGY), 29 October, 1998 (29.10.98), & AU 9871211 A1 & EP 975691 A1 & JP 2002-515932 A	9,10

Form PCT/ISA/210 (continuation of second sheet) (July 1998)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/12091

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BOHL, K. S. et al., Nitric oxide-generating polymers reduce platelet adhesion and smooth muscle cell proliferation. Biomaterials, 2000, Vol.21, No.22, pages 2273 to 2278	9,10

Form PCT/ISA/210 (continuation of second sheet) (July 1998)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/12091

**Box I Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 3 of first sheet)**This International Searching Authority found multiple inventions in this international application, as follows:  
(See extra sheet.)

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.  
☒ No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1998)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP02/12091

Continuation of Box No.II of continuation of first sheet(1)

(1) The special technical feature of the inventions as set forth in claims 1 to 6, 12 and the parts of the inventions as set forth in claims 13 and 14 directly or indirectly depending on the invention as set forth in claim 12 resides in "a drug for preventing vascular restenosis which contains as the active ingredient a compound inactivating  $\alpha$ 1-portease inhibitor".

In contrast, (2) the special technical feature of the inventions as set forth in claims 7 to 11 and the parts of the inventions as set forth in claims 13 and 14 directly or indirectly depending on the invention as set forth in claim 7 resides in "a drug for preventing vascular restenosis which contains as the active ingredient a compound inactivating  $\alpha$ -2 macroglobulin".

Since the compound inactivating  $\alpha$ 1-portease inhibitor and the compound inactivating  $\alpha$ -2 macroglobulin are different from each other, it cannot be said that there is a technical relationship between these groups of inventions (1) and (2) involving one or more of the same or corresponding special technical features. Such being the case, these two groups of inventions are not considered as relating to a group of inventions so linked as to form a single general inventive concept.

Thus, the present application has two groups of inventions.

Form PCT/ISA/210 (extra sheet) (July 1998)